



Semiparametric Estimation of the Intensity of Long Memory in Conditional Heteroskedasticity^{*}

LIUDAS GIRAITIS¹, PIOTR KOKOSZKA², REMIGIJUS LEIPUS³ and
GILLES TEYSSIÈRE⁴

¹*Department of Economics, London School of Economics, Houghton Street, London WC2A 2AE, United Kingdom. e-mail: l.giraitis@lse.ac.uk*

²*Department of Mathematical Sciences, University of Liverpool, Liverpool, L69 3BX, United Kingdom. e-mail: P.S.Kokoszka@liverpool.ac.uk*

³*Department of Mathematics and Informatics, Vilnius University, Naugarduko 24, Vilnius 2600, Lithuania. e-mail: Remigijus.Leipus@Maf.Vu.lt*

⁴*Gilles Teyssière, European Commission, and GREQAM Joint Research Centre, ISIS, TP 361, Via E. Fermi, 1. I-21020 Ispra (VA), Italy. e-mail: gilles.teyssiere@jrc.it; gilles@ehess.cnrs-mrs.fr*

Abstract. The paper is concerned with the estimation of the long memory parameter in a conditionally heteroskedastic model proposed by Giraitis et al. (1999b). We consider estimation methods based on the partial sums of the squared observations, which are similar in spirit to the classical *R/S* analysis, as well as spectral domain approximate maximum likelihood estimators. We review relevant theoretical results and present an empirical simulation study.

Key words: long memory, ARCH models, semiparametric estimation, modified *R/S*, KPSS and *V/S* statistics, periodogram.

1. Introduction

Long memory, a term commonly used to describe persistent dependence between time series observations as the lag increases, has been shown to be present in geophysical and, more recently, in network traffic data. It is, however, still a matter of debate if market data also exhibit some form of long memory. Many earlier studies, focused on the returns themselves. Long memory in returns, or levels, as it is also commonly referred to, would, however, be a radical departure from the random walk hypothesis and the assumption of the unpredictability of asset returns which underlines the classical asset pricing theory. Empirical studies also suggest that the returns are essentially uncorrelated and the presence of a weak correlation can be to a large extent explained by factors like bid-ask spread and non-synchronous trading, see Campbell et al. (1997). However, the presence of strong dependence between the squares or absolute values of returns does not contradict the efficient

^{*} Research partially supported by EPSRC grant GR/L/78222 at the University of Liverpool. Liudas Giraitis was supported by ESRC grant R000235892 and ESRC grant R000238212. Remigijus Leipus was partially supported by Lithuanian Science and Studies Foundation Grant K-014.

market hypothesis and many empirical studies suggest that such transformations of returns exhibit some form of persistent dependence. The presence of long memory in the squares of returns may have profound implications. For example, the volatility estimators based on historical data can be affected, which may in turn impact pricing of derivative products.

In order to develop estimation procedures, a parametric or semiparametric model must be postulated in which the squares of returns form a long memory stationary sequence. Even though several attempts have been made to construct such models by modifying classical ARCH or GARCH specifications, Giraitis et al. (2000) showed that some of these models have in fact short memory, see Section 2 for more details. Recall that in the context of covariance stationary linear time series, long memory is typically characterized by the requirement that the autocovariance function decays at the rate k^{2d-1} , $0 < d < 1/2$, and hence, is not absolutely summable; a series is said to have short memory if the autocovariance function is absolutely summable. These definitions are applicable to any stationary sequences, and we adopt them in this paper to the sequences of squares r_t^2 , where the r_t follow an ARCH type model developed by Giraitis et al. (1999b). The new model is different from the traditional ARCH(∞) in that the parameter σ_t itself, not the conditional variance σ_t^2 , is a linear function of the past returns. The construction implies that the autocovariance function $\text{Cov}(r_t^2, r_{t+k}^2)$ decays at the rate k^{2d-1} for some $0 < d < 1/2$. We believe that it is not possible to modify the classical ARCH(∞) specification in such a way that the autocovariances of the r_t^2 decay like k^{2d-1} , see Proposition 2.1 and Giraitis et al. (1999b) for a more extensive discussion. The model of Giraitis et al. (2000) is described in detail in Section 2.

The paper examines two types of estimation procedures. The first class of estimators, examined from the theoretical and empirical point of view, goes back to the pioneering work of Mandelbrot and his collaborators, see references in Section 3, who developed the *rescaled range*, or R/S , method of Hurst (1951) into a widely used tool for estimating the intensity of long memory. In addition to the R/S method, we also study estimators based on the KPSS statistic of Kwiatkowski et al. (1992) and the V/S statistic proposed by Giraitis et al. (1999a). In the latter two methods, the range of the partial sums appearing in the R/S statistic is replaced, respectively, by their ‘second moment’ and ‘variance’. Details are presented in Subsection 3.1. The above three methods are based on subdividing the sample into a number of blocks. The choice of the blocks is important as it affects the accuracy of the estimators. There is no theoretical guidance as to how to subdivide the sample, so Monte Carlo simulations must be employed. The second procedure is based on the spectral domain approximate maximum likelihood estimator developed by Robinson (1995) in the setting of linear long memory processes. We focus only on an empirical study of this estimator. In a practical implementation of this procedure, the choice of a bandwidth of Fourier frequencies around zero is crucial. Even though some theoretical results are available in the linear and

Gaussian cases, see Subsection 3.2, Monte Carlo simulations offer a more detailed guidance. Our simulations support the conjecture that this estimation procedure, developed originally for Gaussian and linear processes, might also be applicable to ARCH type models. Its theoretical justification and properties in this setting are largely an interesting open problem.

The paper is organized as follows: Section 2 introduces the model of Giraitis et al. (1999b). In Section 3, we describe the estimators and develop the necessary theoretical background. Section 4 contains the results of an extensive simulation study and provides the technical details of the implementation of the estimation procedures presented in Section 3.

2. The Model

We describe in this section the model of Giraitis et al. (1999b) and discuss its main properties. The central feature of this model is that while the observations (returns) r_t are uncorrelated, their squares have the autocovariance function which is not absolutely summable. This is in contrast to a classical ARCH(∞) sequence whose squares have an absolutely summable autocovariance function. To underline the differences between the two specifications, we begin by recalling some relevant properties of the classical ARCH(∞) model.

A random sequence $\{r_k, k \in \mathbf{Z}\}$ is said to satisfy ARCH(∞) equations if there exists a sequence of independent identically distributed zero mean random variables $\{\varepsilon_k, k \in \mathbf{Z}\}$ such that

$$r_k = \sigma_k \varepsilon_k, \quad \sigma_k^2 = a + \sum_{j=1}^{\infty} b_j r_{k-j}^2, \quad (2.1)$$

where $a \geq 0$, $b_j \geq 0$, $j = 1, 2, \dots$. As mentioned in the introduction, in this paper we focus on the sequence of squares $X_k = r_k^2$. If the r_k obey (2.1), then the X_k satisfy the equations

$$X_k = \varepsilon_k^2 \left(a + \sum_{j=1}^{\infty} b_j X_{k-j} \right). \quad (2.2)$$

Using a Volterra-type representation

$$X_k = a + a \sum_{l=1}^{\infty} \sum_{j_1, \dots, j_l=1}^{\infty} b_{j_1} \dots b_{j_l} \varepsilon_k^2 \varepsilon_{k-j_1}^2 \dots \varepsilon_{k-j_1-\dots-j_l}^2,$$

Giraitis et al. (2000) obtained a number of results which show that under mild assumptions sequences X_k satisfying (2.2) cannot have long memory. These assumptions require essentially that $\sum_{j=1}^{\infty} b_j < \infty$, a condition imposed also in Ding and Granger (1996), Baillie et al. (1996) and related papers which aimed

at constructing ARCH type models with long memory in squares. Kokoszka and Leipus (2000) showed that under the assumption

$$(E\varepsilon_0^4)^{1/2} \sum_{j=1}^{\infty} b_j < 1 \quad (2.3)$$

there exists a unique weakly stationary solution to (2.2). Giraitis et al. (1999a, 2000) established the following results which show that the classical ARCH(∞) model (2.1) has short memory in squares.

PROPOSITION 2.1. *If assumption (2.3) is satisfied, then for any $k \in \mathbf{Z}$*

$$0 \leq \text{Cov}(X_k, X_0) < \infty$$

and

$$\sum_{k=-\infty}^{\infty} \text{Cov}(X_k, X_0) < \infty. \quad (2.4)$$

THEOREM 2.1. *Suppose $E\varepsilon_0^8 < \infty$ and*

$$(E\varepsilon_0^8)^{1/4} \sum_{j=1}^{\infty} b_j < 1.$$

Then as $N \rightarrow \infty$

$$N^{-1/2} \sum_{j=1}^{\lfloor Nt \rfloor} (X_j - EX_j) \xrightarrow{D[0,1]} \sigma W(t), \quad (2.5)$$

where $\{W(t), t \in [0, 1]\}$ is the standard Brownian motion, $\xrightarrow{D[0,1]}$ means weak convergence in the space $D[0, 1]$ endowed with the Skorokhod topology and $\sigma^2 = \sum_{k=-\infty}^{\infty} \text{Cov}(X_k, X_0)$.

In a model introduced by Robinson (1991) and developed by Giraitis et al. (1999b), which we call here LM ARCH(∞) (not to be confused with the LM-ARCH of Ding and Granger (1996) which is of the form (2.1)), relations (2.4) and (2.5) no longer hold; the covariances of the X_k decay at the rate k^{2d-1} for some $0 < d < 1/2$, and appropriately normalized partial sums converge to a fractional Brownian motion. This model is defined as follows. The r_k are assumed to satisfy

$$r_k = \sigma_k \varepsilon_k, \quad \sigma_k = \alpha + \sum_{j=1}^{\infty} \beta_j r_{k-j}, \quad (2.6)$$

where $\{\varepsilon_k, k \in \mathbf{Z}\}$ is a sequence of zero mean finite variance i.i.d. random variables, α is a real number and the weights β_j satisfy

$$\beta_j \sim c j^{d-1}, \quad 0 < d < 1/2, \quad (2.7)$$

for some $c > 0$.

Note that neither α nor the β_j are assumed positive and, unlike in (2.1), σ_k , not its square, is a linear combination of the past r_k , rather than their squares. Observe also that condition (2.7) implies only $\sum_j \beta_j^2 < \infty$ which contrasts with the assumption $\sum_j b_j < \infty$.

Giraitis et al. (1999b) established the following results which show that the squares of the r_k satisfying (2.6) and (2.7) have two essential features of long memory: hyperbolically decaying non-summable covariances and attraction to a fractional Brownian motion.

THEOREM 2.2. *Suppose $E\varepsilon_0^4 < \infty$ and*

$$L(E\varepsilon_0^4)^{1/2} \sum_{j=1}^{\infty} \beta_j^2 < 1, \quad (2.8)$$

where $L = 7$ if the ε_k are Gaussian and $L = 11$ in other cases. Then there is a stationary solution to Equations (2.6) and (2.7) given by orthogonal Volterra series

$$r_k = \sigma_k \varepsilon_k, \quad \sigma_k = \alpha \sum_{l=0}^{\infty} \sum_{j_1, \dots, j_l=1}^{\infty} \beta_{j_1} \cdots \beta_{j_l} \varepsilon_{k-j_1} \cdots \varepsilon_{k-j_1-\dots-j_l}. \quad (2.9)$$

The sequence $X_k = r_k^2$ is covariance stationary and as $k \rightarrow \infty$

$$\text{Cov}(X_k, X_0) \sim C k^{2d-1}, \quad (2.10)$$

where C is a positive constant.

Recall that a Gaussian process $\{W_H(t), t \geq 0\}$ is a fractional Brownian motion with parameter $H \in (0, 1)$ if it has mean zero and covariances

$$E[W_H(t_1)W_H(t_2)] = \frac{1}{2}(t_1^{2H} + t_2^{2H} - |t_1 - t_2|^{2H}). \quad (2.11)$$

THEOREM 2.3. *If conditions of Theorem 2.2 are satisfied then as $N \rightarrow \infty$*

$$\frac{1}{N^{1/2+d}} \sum_{j=1}^{[Nt]} (X_j - EX_j) \xrightarrow{D[0,1]} c_d W_{1/2+d}(t), \quad (2.12)$$

where c_d is a positive constant.

We see that, in spite of the nonlinearity, the LM ARCH(∞) model satisfies, up to a scaling constant, the same asymptotic relations (2.10) and (2.12) as the moving average $Y_k = \sum_{j=0}^{\infty} \beta_{k-j} \varepsilon_j$ with the weights β_j (2.7).

We conclude this section by noting that the smallest possible value of L in (2.8) is not known; this is a complex combinatorial problem. In the Gaussian case the third order cumulants in a diagram formula used in the proof vanish, so a smaller value of L can be taken.

In the simulations presented in Section 4 we also use coefficients β_j for which relation (2.8) fails to hold with $L = 7$, so, strictly speaking, there is no theoretical justification for the results obtained in such cases. The estimation procedures, however, continue to perform quite well, suggesting a need for further theoretical research in this direction.

3. The Estimators

In this section, we describe the two estimation procedures for the long memory parameter d in (2.6) and (2.7), and provide some theoretical background. Throughout the present section X_1, \dots, X_N is an observed sample.

3.1. ESTIMATORS BASED ON THE PARTIAL SUMS

We present here a theoretical background for three procedures based on Theorem 2.3. We begin with the rescaled range, or R/S analysis introduced by Hurst (1951) and subsequently refined by Mandelbrot and his collaborators, see Mandelbrot and Wallis (1969), Mandelbrot (1972, 1975) and Mandelbrot and Taqqu (1979).

The R/S statistic is defined as \hat{R}_N/\hat{s}_N where

$$\hat{R}_N = \max_{1 \leq k \leq N} \sum_{j=1}^k (X_j - \bar{X}_N) - \min_{1 \leq k \leq N} \sum_{j=1}^k (X_j - \bar{X}_N) \quad (3.1)$$

is the range and

$$\hat{s}_N^2 = \frac{1}{N} \sum_{j=1}^N (X_j - \bar{X}_N)^2 \quad (3.2)$$

is a standard variance estimator. In (3.1) and (3.2), \bar{X}_N is the sample mean $N^{-1} \sum_{j=1}^N X_j$. The identity

$$\sum_{j=1}^k (X_j - \bar{X}_N) = \sum_{j=1}^k (X_j - EX_j) - \frac{k}{N} \sum_{j=1}^N (X_j - EX_j)$$

and Theorem 2.3 imply that

$$\frac{\hat{R}_N}{N^{1/2+d}} \xrightarrow{d} c_d \left\{ \max_{0 \leq t \leq 1} W_{1/2+d}^0(t) - \min_{0 \leq t \leq 1} W_{1/2+d}^0(t) \right\}, \quad (3.3)$$

where

$$W_{1/2+d}^0(t) = W_{1/2+d}(t) - tW_{1/2+d}(1)$$

is a fractional Brownian bridge, cf. (2.11). It is equally easy to verify that

$$\hat{s}_N^2 \xrightarrow{P} \text{Var}X_1. \quad (3.4)$$

Indeed,

$$\hat{s}_N^2 = \frac{1}{N} \sum_{j=1}^N (X_j^2 - EX_j^2) + (EX_1^2 - [\bar{X}_N]^2). \quad (3.5)$$

By the Volterra representation (2.9) X_j^2 can be written as $X_j^2 = f(\varepsilon_j, \varepsilon_{j-1}, \dots)$ where f is a measurable function. Since $\{\varepsilon_j\}$ is an ergodic sequence this implies (cf. Theorem 3.5.8 of Stout, 1974) ergodicity of $\{X_j^2\}$. Under assumptions of Theorem 2.3 $EX_j^2 < \infty$. Therefore the first term in (3.5) tends to zero. By the same argument as above $\{X_j\}$ is ergodic as well, and therefore $\bar{X}_N \Rightarrow EX_1$. Hence the second term in (3.5) tends to $\text{Var}X_1$.

Combining (3.3) and (3.4), we see that as $N \rightarrow \infty$

$$\frac{1}{N^{1/2+d}} \frac{\hat{R}_N}{\hat{s}_N} \xrightarrow{d} \frac{c_d \{\max_{0 \leq t \leq 1} W_{1/2+d}^0(t) - \min_{0 \leq t \leq 1} W_{1/2+d}^0(t)\}}{(\text{Var}X_1)^{1/2}} =: R_d. \quad (3.6)$$

Relation (3.6) forms a theoretical foundation for the R/S method. Taking logarithms of both sides, we obtain a heuristic identity

$$\log \left(\frac{\hat{R}_N}{\hat{s}_N} \right) \approx \left(\frac{1}{2} + d \right) \log N + \log R_d, \quad \text{as } n \rightarrow \infty,$$

which can also be written as

$$\hat{d}_{R/S} - d = O_P \left(\frac{1}{\log N} \right) \quad \text{with } \hat{d}_{R/S} = \frac{\log(\hat{R}_N/\hat{s}_N)}{\log N} - \frac{1}{2},$$

and which shows that $1/2 + d$ can be interpreted as the slope of a regression line of $\log(\hat{R}_N/\hat{s}_N)$ on $\log N$ with random intercept $\log R_d$. The point of the R/S analysis is to consider many subsamples of varying size N from a given sample X_1, \dots, X_N in order to obtain many points which are used to estimate the slope of the regression line, see for example Mandelbrot and Taqqu (1979) or Beran (1994). The technical details of the implementation of this procedure are described in Section 4.

The above discussion shows that in place of the range (3.1), any other ‘simple’ continuous functional of the partial sum process can form a basis for an estimation procedure of the type just described. We focus below on the KPSS and

V/S statistics, used by Giraitis et al. (1999a) to test for long memory in ARCH models, which provide the other two estimators of d .

The KPSS statistic was introduced by Kwiatkowski et al. (1992) to test trend stationarity against a unit root alternative. Lee and Schmidt (1996) used the KPSS statistic to test for the presence of long memory in a stationary linear time series and gave its asymptotic distribution under long memory alternatives, but provided only heuristic outlines of the proofs.

In the context of testing for long memory in a stationary sequence the KPSS statistic takes the form:

$$\hat{T}_N = \frac{\hat{M}_N}{N\hat{s}_N^2} \quad (3.7)$$

with \hat{s}_N^2 given by (3.2) and

$$\hat{M}_N = \frac{1}{N} \sum_{k=1}^N \left[\sum_{j=1}^k (X_j - \bar{X}_N) \right]^2.$$

We thus see that the range has been replaced by the second moment. We retained the N in the denominator of the RHS of (3.7) in order to conform to the original definition of Lee and Schmidt (1996); unlike the R/S statistic which must be divided by \sqrt{N} in order to ensure convergence for weakly dependent X_j , the statistic \hat{T}_N converges in this case without any normalization.

By Theorem 2.3,

$$\frac{\hat{M}_N}{N^{1+2d}} \xrightarrow{d} c_d^2 \int_0^1 [W_{1/2+d}^0(t)]^2 dt. \quad (3.8)$$

Hence setting $\hat{d}_{\text{KPSS}} = \log \hat{T}_N / (2 \log N)$ we get

$$\hat{d}_{\text{KPSS}} - d = O_P\left(\frac{1}{\log N}\right).$$

Combining relation (3.8) with (3.4), we see that the slope of the regression line of $\log \hat{T}_N$ on $\log N$ estimates $2d$, whereas the regression of $\log(\hat{M}_N^{1/2}/\hat{s}_N)$ on $\log N$ yields an estimate of $d + 1/2$.

In the context of long memory hypothesis testing, Giraitis *et al.* (1999a) proposed the statistic

$$\hat{U}_N = \frac{\hat{V}_N}{\hat{s}_N^2 N}, \quad (3.9)$$

where

$$\hat{V}_N = \frac{1}{N} \left\{ \sum_{k=1}^N \left[\sum_{j=1}^k (X_j - \bar{X}_N) \right]^2 - \frac{1}{N} \left[\sum_{k=1}^N \sum_{j=1}^k (X_j - \bar{X}_N) \right]^2 \right\}.$$

They called \hat{U}_N the V/S statistic for ‘variance over S ’. This statistic is very similar to the KPSS statistic, the second sample moment \hat{M}_N in (3.7) is replaced by the sample variance \hat{V}_N . The statistic \hat{U}_N contains a correction for a mean and is more sensitive to ‘shifts in variance’ than \hat{T}_N , see Giraitis et al. (1999a) for further background and discussion.

Arguing as above, we conclude that the regressions of $\log \hat{U}_N$ and $\log(\hat{V}_N^{1/2}/\hat{s}_N)$ on $\log N$ will, respectively, yield estimates of $2d$ and $d + 1/2$ (setting $\hat{d}_{V/S} = \log \hat{U}_N / (2 \log N)$ we get $\hat{d}_{V/S} - d = O_P(1/\log N)$).

3.2. SPECTRAL DOMAIN ESTIMATION

We describe here the local Whittle estimator proposed by Künsch (1987) and developed by Robinson (1995) which is used to estimate the parameters $C > 0$ and $0 < d < 1/2$ assuming that the observed Gaussian or moving average series has spectral density $f(\lambda)$ which behaves at low frequencies like

$$f(\lambda) \sim C|\lambda|^{-2d}, \quad \lambda \rightarrow 0. \quad (3.10)$$

The estimator minimizes an approximate Gaussian maximum likelihood function:

$$\frac{1}{m} \sum_{j=1}^m \left\{ \ln(C\lambda_j^{-2d}) + \frac{I(\lambda_j)}{C\lambda_j^{-2d}} \right\},$$

where

$$I(\lambda_j) = \frac{1}{2\pi N} \left| \sum_{k=1}^N X_k e^{ik\lambda_j} \right|^2$$

is the periodogram at the Fourier frequencies $\lambda_j = 2\pi j/N$, $j = 1, \dots, m$. The bandwidth m increases more slowly than the sample size N :

$$\frac{1}{m} + \frac{m}{N} \rightarrow 0 \quad \text{as } N \rightarrow \infty.$$

Robinson (1995) showed that under appropriate conditions, which include the existence of a linear moving average representation, the estimator of d is asymptotically normal and converges at the rate \sqrt{m} :

$$\sqrt{m}(\hat{d} - d) \sim \mathcal{N}(0, \frac{1}{4}),$$

where the upper bound for m depends on the degree of the smoothness of spectral density $f(\lambda)$ when $\lambda \rightarrow 0$.

In the case of long-memory LM ARCH(∞) sequences discussed in Theorem 2.2, no similar asymptotic theory is available at present. However, note that relation (2.10) implies that the spectral density f of the sequence $X_k = r_k^2$ satisfies (3.10).

Thus, although the local Whittle estimator was designed for Gaussian or moving average time series, we expect that it is applicable also to the LM ARCH(∞) series with the Volterra representation (2.9). This is because the weights β_1, β_2, \dots can be conveniently factorized and are square summable. We expect that, similarly as for moving averages, these properties effectively control the dependence structure of the X_k and allow to derive not only the CLT, Theorem 2.2, but also the asymptotic distribution of the local Whittle estimator. The empirical results show that this is at least a consistent estimator of d .

In the Gaussian case, the problem of the choice of the bandwidth m is related to the smoothness of the short memory component $h(\lambda)$ appearing in the following factorization of the spectral density:

$$f(\lambda) = |1 - \exp(i\lambda)|^{-2d} h(\lambda).$$

Assuming that h is twice differentiable and $h(0) > 0$, Delgado and Robinson (1996) proved that in case of the modified Geweke Porter-Hudak estimator the optimal m is given by

$$m_{\text{optimal}} = c_{\text{optimal}} N^{4/5}, \quad (3.11)$$

where

$$c_{\text{optimal}} = \left(\frac{3}{4\pi} \right)^{4/5} \left(\frac{h''(0)}{2h(0)} + \frac{1}{12}d \right)^{-2/5}. \quad (3.12)$$

The use of (3.11) and (3.12) in the case of the local Whittle estimator is not theoretically justified and not fully convincing even for Gaussian sequences, let alone for the LM ARCH(∞) model. Nevertheless, we use these formulas in our setting and evaluate the optimal bandwidth from the data by using an iterative procedure proposed by Robinson and Henry (1996). We evaluate the quality of an estimator constructed in this way by comparing it with the R/S type estimators described in Subsection 3.1.

4. Simulations

We consider the following data generating process (DGP):

$$r_k = \sigma_k \varepsilon_k, \quad \sigma_k = \alpha + \sum_{j=1}^{\infty} \beta_j r_{k-j}, \quad k = 1, \dots, N, \quad \varepsilon_k \sim N(0, 1). \quad (4.13)$$

Two sample sizes are considered, $N = 3000, 6000$. To reduce initialization effects, which are particularly strong for long-range dependent sequences, we generate for each simulation a series $r_{-\tau}, r_{-\tau+1}, \dots, r_{-1}, r_0, r_1, \dots, r_N$, $\tau = 12000$, where the pre-sample observations $r_{-\tau}, r_{-\tau+1}, \dots, r_{-1}, r_0$ are recursively used for initializing the process, and are discarded afterwards. We truncate the infinite order lag polynomial $\beta(L)$ at the order 5000 to take into account the dependence between

Table I. Estimation results for the LM ARCH(∞) process, Model A (root mean squared error in parentheses)

d	3000 observations				6000 observations			
	V/S	R/S	KPSS	Robinson	V/S	R/S	KPSS	Robinson
0.05	0.0092 (0.0622)	0.0495 (0.0362)	0.0114 (0.0662)	0.0285 (0.0362)	0.0122 (0.0527)	0.0479 (0.0284)	0.0137 (0.0557)	0.0294 (0.0307)
0.075	0.0321 (0.0645)	0.0677 (0.0378)	0.0363 (0.0674)	0.0553 (0.0378)	0.0371 (0.0536)	0.0682 (0.0301)	0.0406 (0.0556)	0.0560 (0.0324)
0.1	0.0800 (0.0528)	0.1102 (0.0391)	0.0835 (0.0585)	0.0834 (0.0396)	0.0826 (0.0425)	0.1093 (0.0317)	0.0848 (0.0470)	0.0852 (0.0326)
0.125	0.0949 (0.0591)	0.1190 (0.0398)	0.1033 (0.0620)	0.1125 (0.0413)	0.1035 (0.0460)	0.1235 (0.0319)	0.1103 (0.0487)	0.1145 (0.0331)
0.15	0.1284 (0.0562)	0.1470 (0.0406)	0.1383 (0.0604)	0.1395 (0.0437)	0.1375 (0.0438)	0.1527 (0.0333)	0.1454 (0.0480)	0.1432 (0.0345)
0.175	0.1599 (0.0550)	0.1738 (0.0415)	0.1708 (0.0606)	0.1656 (0.0471)	0.1689 (0.0429)	0.1801 (0.0343)	0.1775 (0.0485)	0.1708 (0.0370)
0.2	0.1776 (0.0578)	0.1887 (0.0434)	0.1894 (0.0619)	0.1701 (0.0579)	0.1872 (0.0448)	0.1959 (0.0347)	0.1965 (0.0490)	0.1811 (0.0439)
0.225	0.1668 (0.0790)	0.1786 (0.0625)	0.1801 (0.0761)	0.1778 (0.0688)	0.1787 (0.0632)	0.1874 (0.0510)	0.1898 (0.0604)	0.1902 (0.0531)

very distant observations. The sequence of innovations ε_k is generated by using two different random number generators, with different seeds. We randomly draw one of the two generated uniform deviates, and transform it to a Gaussian random variable by using the Box-Muller method.

We generate 5000 independent samples. Each sample of N observation is subdivided into B adjacent and non-overlapping blocks of observations of equal size $[N/B]$. We then obtain a grid $t_1 = 1, t_2 = [N/B] + 1, \dots, t_i = (i - 1)[N/B] + 1, \dots, t_B = N - [N/B] + 1$. For each point of the sequence $\{t_i\}_{i=1}^B$ we define a sequence of K increasing nested blocks with origin t_i , i.e., $\{[t_i, t_i + k_j]\}_{j=1}^K$, such that $t_i + k_j \leq N$, the sequence of K steps $\{k_j\}_{j=1}^K$ is given by a logarithmic grid. Interested readers are referred to Beran (1994), p. 84–85, for more details on the ‘pox-plot’ based estimators. Beran (1994) reports the fact that the ‘pox-plot’ based estimates of the Hurst exponent of the Nile river data strongly depend on the choice of K , and that “it seems difficult to define an ‘automatic’ ‘pox-plot’ methodology, and to derive results on statistical inference based on the method”. In our simulations, the minimum value of K is set to 40 and the number of blocks B is set to 40. This choice is motivated by the simulation results.

We calculate the R/S , V/S and KPSS statistics for each interval $\{[t_i, t_i + k_j]\}_{i=1}^B\}_{j=1}^K$ and obtain the sequences $\{\{R/S(t_i, k_j)\}_{i=1}^B\}_{j=1}^K$, $\{\{V/S(t_i, k_j)\}_{i=1}^B\}_{j=1}^K$, and $\{\{KPSS(t_i, k_j)\}_{i=1}^B\}_{j=1}^K$. We plot the logarithm of the statistics $\log(R/S(t_i, k_j))$,

Table II. Estimation results for the LM ARCH(∞) process, Model B (root mean squared error in parentheses)

d	3000 observations				6000 observations			
	V/S	R/S	KPSS	Robinson	V/S	R/S	KPSS	Robinson
0.05	0.0048 (0.0643)	0.0434 (0.0358)	0.0069 (0.0678)	0.0138 (0.0470)	0.0075 (0.0556)	0.0414 (0.0290)	0.0085 (0.0584)	0.0153 (0.0417)
0.075	0.0041 (0.0846)	0.0453 (0.0462)	0.0066 (0.0865)	0.0069 (0.0742)	0.0085 (0.0756)	0.0449 (0.0410)	0.0105 (0.0767)	0.0078 (0.0708)
0.1	0.0227 (0.0907)	0.0597 (0.0543)	0.0276 (0.0907)	0.0213 (0.0845)	0.0297 (0.0797)	0.0615 (0.0482)	0.0342 (0.0788)	0.0244 (0.0794)
0.125	0.0502 (0.0894)	0.0815 (0.0576)	0.0582 (0.0875)	0.0446 (0.0877)	0.0603 (0.0756)	0.0861 (0.0493)	0.0675 (0.0731)	0.0516 (0.0788)
0.15	0.0841 (0.0833)	0.1088 (0.0569)	0.0949 (0.0804)	0.0764 (0.0837)	0.0963 (0.0673)	0.1158 (0.0467)	0.1059 (0.0644)	0.0873 (0.0705)
0.175	0.1208 (0.0755)	0.1390 (0.0543)	0.1338 (0.0731)	0.1133 (0.0755)	0.1342 (0.0585)	0.1478 (0.0429)	0.1454 (0.0566)	0.1268 (0.0593)
0.2	0.1573 (0.0686)	0.1697 (0.0517)	0.1718 (0.0680)	0.1512 (0.0673)	0.1709 (0.0518)	0.1796 (0.0398)	0.1831 (0.0520)	0.1662 (0.0501)
0.225	0.1914 (0.0639)	0.1989 (0.0501)	0.2067 (0.0653)	0.1875 (0.0620)	0.2046 (0.0480)	0.2093 (0.0383)	0.2171 (0.0503)	0.2032 (0.0450)
0.25	0.2215 (0.0615)	0.2251 (0.0498)	0.2370 (0.0642)	0.2204 (0.0607)	0.2340 (0.0465)	0.2357 (0.0381)	0.2464 (0.0501)	0.2364 (0.0444)
0.275	0.2468 (0.0612)	0.2474 (0.0512)	0.2622 (0.0640)	0.2490 (0.0629)	0.2584 (0.0465)	0.2579 (0.0393)	0.2703 (0.0500)	0.2647 (0.0472)
0.3	0.2676 (0.0624)	0.2661 (0.0544)	0.2828 (0.0641)	0.2728 (0.0663)	0.2766 (0.0491)	0.2750 (0.0434)	0.2876 (0.0510)	0.2892 (0.0533)
0.325	0.2831 (0.0659)	0.2799 (0.0609)	0.2985 (0.0648)	0.2909 (0.0731)	0.2887 (0.0576)	0.2864 (0.0533)	0.2988 (0.0575)	0.3048 (0.0611)
0.35	0.2918 (0.0774)	0.2880 (0.0744)	0.3057 (0.0742)	0.3038 (0.0843)	0.2968 (0.0687)	0.2943 (0.0664)	0.3061 (0.0664)	0.3176 (0.0736)
0.375	0.2963 (0.0932)	0.2922 (0.0923)	0.3095 (0.0879)	0.3115 (0.0992)	0.2998 (0.0865)	0.2976 (0.0853)	0.3083 (0.0827)	0.3238 (0.0896)

$\log(V/S(t_i, k_j))$, $\log(\text{KPSS}(t_i, k_j))$, against $\log(k_j)$ and then obtain a ‘pox-plot’. Let \hat{b} be the slope of the least-squares regression line fitted to these pox-plots. Then $\hat{d}_{R/S} = \hat{b} - 1/2$, $\hat{d}_{V/S} = \hat{b}/2$, and $\hat{d}_{\text{KPSS}} = \hat{b}/2$.

We consider three DGP-s which differ by the parameterization of the infinite order lag polynomial $\beta(L)$: we have chosen three parameterizations of the moving average form of a FARIMA process. For all models, the parameter α , which is the initial value of the process, is set to 0.10.

Table III. Estimation results for the LM ARCH(∞) process, Model C (root mean squared error in parentheses)

d	3000 observations				6000 observations			
	V/S	R/S	KPSS	Robinson	V/S	R/S	KPSS	Robinson
0.05	-0.0064 (0.0723)	0.0374 (0.0370)	-0.0058 (0.0760)	-0.0023 (0.0600)	-0.0053 (0.0656)	0.0344 (0.0316)	-0.0053 (0.0686)	-0.0023 (0.0568)
0.075	-0.0001 (0.0879)	0.0421 (0.0482)	0.0020 (0.0900)	0.0028 (0.0779)	0.0033 (0.0802)	0.0408 (0.0441)	0.0049 (0.0814)	0.0035 (0.0748)
0.1	0.0160 (0.0963)	0.0544 (0.0581)	0.0205 (0.0962)	0.0151 (0.0901)	0.0226 (0.0858)	0.0558 (0.0527)	0.0269 (0.0848)	0.0176 (0.0857)
0.125	0.0417 (0.0965)	0.0745 (0.0628)	0.0494 (0.0942)	0.0359 (0.0952)	0.0517 (0.0828)	0.0789 (0.0550)	0.0589 (0.0798)	0.0425 (0.0870)
0.15	0.0748 (0.0906)	0.1009 (0.0626)	0.0856 (0.0869)	0.0662 (0.0923)	0.0874 (0.0744)	0.1081 (0.0524)	0.0972 (0.0704)	0.0772 (0.0794)
0.175	0.1117 (0.0820)	0.1311 (0.0596)	0.1251 (0.0783)	0.1032 (0.0836)	0.1259 (0.0644)	0.1403 (0.0478)	0.1376 (0.0609)	0.1174 (0.0669)
0.2	0.1494 (0.0737)	0.1625 (0.0561)	0.1644 (0.0714)	0.1424 (0.0734)	0.1640 (0.0559)	0.1730 (0.0435)	0.1768 (0.0543)	0.1583 (0.0554)
0.225	0.1851 (0.0675)	0.1930 (0.0534)	0.2011 (0.0673)	0.1803 (0.0658)	0.1994 (0.0505)	0.2041 (0.0407)	0.2127 (0.0513)	0.1974 (0.0474)
0.25	0.2172 (0.0638)	0.2209 (0.0521)	0.2336 (0.0654)	0.2153 (0.0621)	0.2307 (0.0478)	0.2322 (0.0397)	0.2439 (0.0505)	0.2328 (0.0446)
0.275	0.2448 (0.0619)	0.2455 (0.0521)	0.2611 (0.0642)	0.2470 (0.0619)	0.2570 (0.0473)	0.2562 (0.0403)	0.2696 (0.0504)	0.2636 (0.0461)
0.30	0.2679 (0.0628)	0.2660 (0.0549)	0.2840 (0.0645)	0.2724 (0.0643)	0.2763 (0.0510)	0.2743 (0.0450)	0.2879 (0.0526)	0.2893 (0.0517)
0.325	0.2839 (0.0673)	0.2806 (0.0615)	0.2993 (0.0672)	0.2937 (0.0720)	0.2918 (0.0555)	0.2888 (0.0514)	0.3025 (0.0554)	0.3092 (0.0584)
0.35	0.2961 (0.0746)	0.2916 (0.0717)	0.3108 (0.0720)	0.3083 (0.0801)	0.3018 (0.0650)	0.2985 (0.0629)	0.3118 (0.0629)	0.3236 (0.0683)
0.375	0.3024 (0.0883)	0.2975 (0.0876)	0.3164 (0.0834)	0.3181 (0.0933)	0.3066 (0.0806)	0.3035 (0.0799)	0.3158 (0.0769)	0.3320 (0.0823)

- Model A. The coefficients β_j are those of the moving average representation of a FARIMA(0, d , 0) process, that is, the coefficients of the fractional difference operator $(1 - L)^{-d}$. Thus, $\beta_j = b_j$, with the b_j defined recursively by $b_0 = 1$, $b_1 = d$, $b_j = b_{j-1}(j - 1 + d)/j$, $j > 1$.
- Model B. The coefficients of this DGP are those of the MA representation of a FARIMA(1, d , 0) process with AR polynomial equal to $1 - \phi L$, that is, of the filter $(1 - \phi L)^{-1}(1 - L)^{-d}$. Thus, $\beta_1 = b_1 + \phi$, $\beta_j = \sum_{k=0}^j \phi^k b_{j-k}$ $j > 1$.

- Model C. The coefficients of this DGP are those of the MA representation of a FARIMA(0, d , 1) process, the MA polynomial being equal to $1 - \theta L$. Thus, the long memory filter is equal to $(1 - \theta L)(1 - L)^{-d}$ and $\beta_1 = b_1 - \theta$, $\beta_j = b_j - \theta b_{j-1}$ $j > 1$.

For the Model A, condition (2.8) is satisfied if $d < 0.1865$. If this condition is not satisfied, there is a large systematic bias for the ‘pox-plot’ based estimators. For that reason, we do not report the estimates for $d > 0.225$. Condition (2.8) can be satisfied by multiplying all the β_j by a constant < 1 . However, Monte Carlo simulation results show that this rescaling leads to a systematic negative bias, that is, the parameter d is underestimated.

For Models B and C, we choose $\theta = 0.20$ and $\phi = -0.20$. These values ensure that condition (2.8) is satisfied for $d > 0.1865$. The coefficients β_j depend on d , but also on the parameters θ and ϕ . If the first elements of the sequence of the β_j are small, there is a systematic negative bias, that is, the parameter d is underestimated. This negative bias is quite large for small values of d , and becomes smaller when $d \in (0.20, 0.375)$, and increases for $d \geq 0.375$.^{*} It is well-known, see e.g. Mandelbrot and Taqqu (1979), that for linear long-memory time series the R/S estimator overestimates d for small d and underestimates it for d close to 0.5, that is, has a bias toward the central values in the range $(0, 0.5)$. It is an interesting finding of our experiments that for the LM ARCH(∞) model, there appears to be a systematic negative bias.

For all the models, it appears that the root mean squared error (RMSE) of the R/S estimator is slightly smaller than the RMSE of the other estimators. The fact that the rate of convergence of the local Whittle estimator is slow and similar to the rate of the other estimators can be a consequence of the choice of the Robinson and Henry (1996) optimal bandwidth which has been developed in the framework of Gaussian long-memory models, with some additional assumptions on the functional form and the smoothness of the spectrum in the neighborhood of the zero frequency. Given that these restrictions might be too strong, some authors, for example, Lobato and Savin (1998), use the local Whittle estimator on a grid of bandwidths.

5. Conclusions

We have considered in this paper several methods for estimating the degree of long-memory in the conditional heteroskedastic model of Giraitis et al. (1999b). Three of these estimators are similar to the ‘pox-plot’ R/S estimator, the fourth one is a spectral domain estimator developed originally for Gaussian time series. Our Monte Carlo simulation results show that these estimators have similar biases and MSE’s which are comparable to those that the R/S method gives for linear time series. Our overall conclusion is that these four estimators, similarly as the

^{*} The whole set of results for $d \in [0.05, 0.5]$ is available upon request from Gilles Teysnière.

R/S analysis for linear time series, can be used as exploratory tools until better estimation procedures become available. Interestingly, unlike for linear models, for the LM ARCH(∞) the bias is always downwards. Another somewhat surprising observation is that in terms of the mean squared error the R/S estimator performs slightly better than the local Whittle estimator; in the Gaussian case the latter has a much better rate of convergence than the R/S estimator, see Robinson (1995) and Giraitis et al. (1999c) among others. This may be due to the lack of an appropriate bandwidth selection procedure for the local Whittle estimator, as discussed towards the end of Subsection 3.2.

The empirical study presented in this paper leaves open many interesting theoretical and practical questions, and we hope that it will stimulate further research.

Acknowledgement

The authors thank the referees for their useful comments.

References

- Baillie, R. T., Bollerslev, T. and Mikkelsen, H. O.: Fractionally integrated generalized autoregressive conditional heteroskedasticity, *J. Economet.* **74** (1996), 3–30.
- Beran, J.: *Statistics for Long-Memory Processes*, Chapman & Hall, New York, 1994.
- Campbell, J. Y., Lo, A. W. and MacKinlay, A. C.: *The Econometrics of Financial Markets*, Princeton University Press, Princeton, 1997.
- Delgado, M. A. and Robinson, P. M.: Optimal spectral bandwidth for long memory, *Statistica Sinica* **6** (1996), 97–112.
- Ding, Z. and Granger, C. W. J.: Modeling volatility persistence of speculative returns: a new approach, *J. Economet.* **73** (1996), 185–215.
- Giraitis, L., Kokoszka, P. and Leipus, R.: Stationary ARCH models: dependence structure and Central Limit Theorem, *Econometric Theory* **16** (2000), 3–22.
- Giraitis, L., Kokoszka, P. and Leipus, R.: Rescaled variance and related tests for long-memory in volatility and levels, Preprint, 1999a.
- Giraitis, L., Robinson, P. and Surgailis, D.: A model for long memory conditional heteroskedasticity, *Ann. Appl. Probab.* 1999b, in press.
- Giraitis, L., Robinson, P. and Surgailis, D.: Variance-type estimation of long memory, *Stochastic Processes and Their Applications* **80**, 1–24.
- Hurst, H.: Long term storage capacity of reservoirs, *Transac. Amer. Soc. Civil Engrs* **116** (1951), 770–799.
- Kokoszka, P. and Leipus, R.: Change-point estimation in ARCH models, *Bernoulli* **6** (2000), in press.
- Künsch, H.: Statistical aspects of self-similar processes, in Yu. A. Prohorov and V. V. Sazonov (eds), *Proceedings of the 1st World Congress of the Bernoulli Society*, vol. 1, VNU Science Press, Utrecht, 1987, pp. 67–74.
- Kwiatkowski, D., Phillips, P. C. B., Schmidt, P. and Shin, Y.: Testing the null hypothesis of stationarity against the alternative of a unit root: how sure are we that economic time series have a unit root? *J. Economet.* **54** (1992), 159–178.
- Lee, D. and Schmidt, P.: On the power of the KPSS test of stationarity against fractionally-integrated alternatives, *J. Economet.* **73** (1996), 285–302.
- Lobato, I and Savin N. E.: Real and spurious long-memory properties of stock market data, *J. Business and Eco. Stat.* **16** (1998), 261–283.

- Mandelbrot, B. B.: Statistical methodology for non-periodic cycles: from the covariance to R/S analysis, *Ann. Eco. Social Measurement* **1** (1972), 259–290.
- Mandelbrot, B. B.: Limit theorems of the self-normalized range for weakly and strongly dependent processes, *Z. Wahrschein. verw. Gebiete* **31** (1975), 271–285.
- Mandelbrot, B. B. and Taqqu, M. S.: Robust R/S analysis of long run serial correlation, in *42nd Session of the International Statistical Institute, Manila*, Book 2, 1979, pp. 69–90.
- Mandelbrot, B. B. and Wallis, J. M.: Robustness of the rescaled range R/S in the measurement of noncyclic long run statistical dependence, *Water Resour. Res.* **5** (1969), 967–988.
- Robinson, P. M.: Testing for strong serial correlation and dynamic conditional heteroskedasticity in multiple regression, *J. Economet.* **47** (1991), 67–84.
- Robinson, P. M.: Gaussian semiparametric estimation of long range dependence, *Ann. Stat.* **23** (1995), 1630–1661.
- Robinson, P. M. and Henry, M.: Bandwidth choice in Gaussian semiparametric estimation of long-range dependence, in P. M. Robinson and M. Rosenblatt (eds), *Athens Conference on Applied Probability and Time Series Analysis*, vol. II: Time Series Analysis. In memory of E. J. Hannan, Springer-Verlag, New York, 1996, pp. 220–232.
- Stout, W. F.: *Almost Sure Convergence*, Academic Press, New York, 1974.